

INTERNATIONAL JOURNAL OF ADVANCES IN  
PHARMACY, BIOLOGY AND CHEMISTRY

Research Article

**Evaluation of the Antiulcer Properties of Castor  
Plants Indigenous to Kerala**

Ashwathy G, Sheela D

Department of Botany, St. Teresa's College, Ernakulam, Kerala, India - 682011.

**ABSTRACT**

The castor oil plant (*Ricinus communis* L.) is a species of flowering plant of Euphorbiaceae family. It is also widely grown as an ornamental plant and for oil seed production in tropical and subtropical regions of the world. The aim of the study was to evaluate the antiulcer properties of the castor oil plant. The antiulcer property was studied by studying the H<sup>+</sup>, K<sup>+</sup>-ATPase enzyme inhibition activity. The H<sup>+</sup>, K<sup>+</sup>-ATPase activity was conducted in the presence of different concentrations of castor seed extract. The percentages of inhibition were calculated. The percentages of inhibition obtained were 94.11%, 94.62%, 94.95%, 95.12% and 95.46% for samples with concentrations 100µl, 200µl, 300µl, 400µl and 500µl respectively. Phytochemical analysis was performed to reveal the presence of antiulcerogenic principles like alkaloids, flavonoids, phenolic compounds and saponins. Positive results were observed for all the four phytochemicals. This showed that *Ricinus communis* L. possess significant antiulcer property.

**Key words:** *Ricinus communis*, antiulcer, H<sup>+</sup>, K<sup>+</sup>-ATPase, percentage of inhibition, phytochemical analysis

**1. INTRODUCTION**

Peptic ulcers are one of the most common disease occurring in human beings. Peptic ulcer disease, also known as a peptic ulcer or stomach ulcer, is a break in the lining of the stomach. It is a chronic and recurrent disease and one of the most predominant of gastrointestinal diseases<sup>1</sup>. Treatment against peptic ulcers includes H<sub>2</sub> antagonists or proton-pump inhibitors, antibiotics or even a surgery<sup>2</sup>. The present study was carried out for the evaluation of the antiulcer properties of castor oil plants indigenous to Kerala. *Ricinus communis* L., the castor oil plant, is a species of flowering plant of Euphorbiaceae family. It is the sole species in the monotypic genus *Ricinus*, and *R. communis* L. is the only species with many polymorphic forms. It is also widely grown as an ornamental plant. It is a tall glabrous glaucous annual shrub. *Ricinus* has been freely used all over India since centuries<sup>3</sup>. The seeds, seed oil, leaves and the roots have great medicinal value. In Indian system of medicine, the leaf, root, and seed oil have been used for the treatment of inflammation and liver diseases as they have been found to be hepatoprotective, laxative, purgative and diuretic<sup>4-8</sup>. Mainly the

seeds and oil of the plant *Ricinus communis* L. has antiulcer properties<sup>4</sup>.

The extract of *Ricinus communis* L. was reported to show antiulcer property in various antiulcer models<sup>8, 9</sup>, but still its activity on H<sup>+</sup>, K<sup>+</sup>-ATPase remains to be unknown. On the basis of these observations, an attempt was made to determine the activity of the castor seed extract on the inhibition of H<sup>+</sup>, K<sup>+</sup>-ATPase enzyme. The inhibition of the H<sup>+</sup>, K<sup>+</sup>-ATPase activity can show whether a plant material has the ability to be an antiulcer agent or not<sup>2, 10-12</sup>. The mechanism of the antiulcer property of *R. communis* L. can thus be studied by studying the H<sup>+</sup>, K<sup>+</sup>-ATPase inhibition activity. H<sup>+</sup>, K<sup>+</sup>-ATPase (Gastric hydrogen potassium ATPase) is an enzyme whose purpose is to acidify the stomach. Excess acidification of the stomach may lead to the formation of peptic ulcers.

The castor plant also possesses antiulcerogenic principles like flavonoids, tannins, alkaloids and saponins<sup>5, 6, 13-17</sup>. Phytochemical analysis can be carried out to find out presence or absence of these antiulcerogenic principles.

## 2. MATERIALS AND METHODS

### 2.1 Plant material:

Fresh twigs of *Ricinus communis* L. were collected from different locations of Ernakulam District, Kerala, India for obtaining the fruit and were identified using the Flora of Presidency of Madras by J. S. Gamble<sup>18</sup>. The thorny coat were peeled and the seeds were collected and air dried.

### 2.2 Preparation of the seed extract:

The seeds collected were grounded into powder. 5g of the powder was weighed and the extract was prepared using chloroform. It was kept undisturbed for 3 days and extract was then filtered and used for the further study.

### 2.3 Preliminary phytochemical analysis:

Phytochemical tests were carried out in order to find out the presence of phytochemical constituents like alkaloids (Dragendroff's Test), flavonoids (Concentrated Sulphuric acid Test), phenolic compounds (Ferric chloride Test), saponins (Foam test) etc. according to the standard procedures<sup>19, 20, 21</sup>.

### 2.4 Analysis of proton pump activity:

#### 2.4.1 Preparation of H<sup>+</sup>, K<sup>+</sup>- ATPase

Gastric membrane containing H<sup>+</sup>, K<sup>+</sup>- ATPase was prepared from mucosal stomach scrapings of sheep and was homogenized in 20mM Tris- HCl buffer (pH 7.4). The homogenate was centrifuged for 20minutes at 15,000 rpm and the resulting supernatant was used to determine the H<sup>+</sup>, K<sup>+</sup>- ATPase activity and its inhibition.

#### 2.4.2 Estimation of protein

The protein content of the supernatant was determined by Lowry's method (1951) using Bovine serum albumin (BSA) as standard.

#### 2.4.3 H<sup>+</sup>, K<sup>+</sup>- ATPase assay

The enzyme extract of 100µl quantity was taken for testing the activity of H<sup>+</sup>, K<sup>+</sup>- ATPase. Reaction was carried out in 16mM Tris- HCl buffer (pH 6.5). The reaction was initiated by adding substrate (2mM ATP, 2mM MgCl<sub>2</sub> and 10mM KCl) and then made upto 2ml and incubated for 30min at 37° C. The reaction was stopped by the addition of an assay mixture containing 45% Ammonium molybdate and 60% Perchloric acid. Phosphomolybdate formed was measured spectrophotometrically at 400nm.

#### 2.4.4 Inhibition of H<sup>+</sup>, K<sup>+</sup>- ATPase in- vitro

The seed extract having different concentrations (100µl, 200µl, 300µl, 400µl, 500µl) were incubated

with H<sup>+</sup>, K<sup>+</sup>- ATPase assay mixture for 30min. O.D values were noted. The results were expressed as percentage of inhibition of enzyme. The percentage of inhibition was calculated using the formula:

Percentage of inhibition =

$$\frac{[O.D_{(control)} - O.D_{(test)}] \times 100}{O.D_{(control)}}$$

O.D<sub>(control)</sub>

## 3. RESULTS

### 3.1 Phytochemical analysis:

The phytochemical screening of *Ricinus communis* L. shows the occurrence of alkaloids, phenolics, flavonoids and saponins (Table 1).

### 3.2 Analysis of proton pump activity:

#### 3.2.1 Estimation of protein

The protein content was found out to be 0.44mg/ml using Lowry's method.

#### 3.2.2 Inhibition of H<sup>+</sup>, K<sup>+</sup>- ATPase in- vitro

The H<sup>+</sup>, K<sup>+</sup>- ATPase inhibitory activity of *Ricinus communis* L. was studied by using different concentrations of the castor seed extract. For the different concentrations of extract taken were 100µl, 200µl, 300µl, 400µl and 500µl, the percentages of inhibition obtained were 94.11%, 94.62%, 94.95%, 95.12% and 95.46% (Table 2). The data showed that the percentage of inhibition increases as the concentration of the extract increases ie, with an increase in the concentration of the extract the antiulcer activity increases.

## 4. DISCUSSION

Castor oil plant, *Ricinus communis* L. has many uses in day-to-day life. Castor oil has been widely reported to be used as a laxative, purgative and cathartic in Unani, Ayurvedic and other ethnomedical systems<sup>4</sup>. In addition to the above mentioned uses, castor plant also has analgesic, antiarthritic, antiinflammatory, antiasthmatic, antidandruff, antinociceptive, antimicrobial, antifungal, hepatoprotective, bioinsecticidal activity, etc<sup>7, 8, 13, 14, 16</sup>. Castor oil plant also possesses significant antiulcerogenic property<sup>4</sup>.

A large number of phytochemicals such as flavonoids, tannins, alkaloids and saponins from other plants have previously demonstrated potential antiulcer activity<sup>8, 10, 22</sup>. In the present study, the presence phytochemical constituents like flavonoids, phenolics, tannins and saponins etc, were found out (Table 1). This proves that *Ricinus communis* L. could be antiulcer agent. The results were similar to works done previously<sup>6, 13, 15, 17, 22, 23</sup>.

The antiulcer property was confirmed by the H<sup>+</sup>, K<sup>+</sup>-ATPase inhibitory activity. The results of the present study showed percentages of inhibition of H<sup>+</sup>, K<sup>+</sup>-ATPase to be more than 90% with different concentrations of the castor seed extract. Therefore, *Ricinus communis* L. proved to be a good plant with significant antiulcer property. Similar studies were carried out in other plants<sup>2, 10, 11, 12</sup>. The Castor oil plant (*Ricinus communis* L.) can thus be used as an antiulcer drug with much reduced side effects as compared to chemical drugs to treat Peptic ulcers after proper purification and formulation.

## 5. CONCLUSION

The present study has shown that Castor oil plant has significant antiulcerogenic property owing to the high H<sup>+</sup>, K<sup>+</sup>-ATPase inhibitory activity as well as to the presence of antiulcerogenic principles like alkaloids, flavonoids, phenolics and saponins. The present study suggests the use of castor oil as a remedy for peptic ulcers after purification and with proper pharmaceutical formulation. Further studies, on the H<sup>+</sup>, K<sup>+</sup>-ATPase inhibitory activity of castor oil are however, recommended to evaluate its antiulcerogenic power.

**Table 1**  
**Result of Phytochemical Analysis**

Phytoconstituents	Castor oil
Alkaloids	+
Phenolics	+
Flavonoids	+
Saponins	+

Key: (+); Present, (-); Absent

**Table 2**  
**Percentage of inhibition of H<sup>+</sup>, K<sup>+</sup>-ATPase enzyme**

Concentration of plant extract (in µl)	Percentage of inhibition (%)
100	94.11
200	94.62
300	94.95
400	95.12
500	95.46

## 6. REFERENCES

1. Kore Kakasaheb, Shete Rajkumar V, Patel Apsari J, Kulkarni Jitendra B. Antiulcer of Aqueous extract of *Spinacia oleracea* in rats. International Journal of Research in Pharmacy and Chemistry, 2011; 1(3): 654-661.
2. Samuel Onasanwo A, Neetu Singh, Samuel Olaleye B, Gautam Palit. Antiulcerogenic and proton pump (H<sup>+</sup>, K<sup>+</sup>-ATPase) inhibitory activity of *Kolaviron* from *Garcinia kola* Haeckel. in rodents. Indian Journal of Experimental Biology, 2011; 49: 461-468.
3. Padma Laxmikant Ladda, Rupali Bhimashankar Kamthane. *Ricinus communis* (Castor): An Overview. International Journal of Research in Pharmacology and Pharmacotherapeutics, 2014; 3 (2): 136-144.
4. Preeti KM, Ansu Bala Verma N. A Review on Ethanopharmacological Potential of *Ricinus communis* Linn.. Pharma Tutor Magazine, 2014; 2(3): 76-85.
5. Gupta Neelam, Singh AK. A Review on *Ricinus communis* Linn., International Ayurvedic Medical Journal, 2015; 3(2): 492-495.
6. Jitendra Jena, Ashish Kumar Gupta. *Ricinus communis* Linn: A Phytopharmacological Review. International Journal of Pharmacy and Pharmaceutical Sciences, 2012; 4(2): 25-29.
7. Vaishali Murade, Dinesh Hase, Keshav Deshmukh, Shreyas Pansambal. A Comprehensive Review of Phytopharmacology of *Ricinus communis*

- (Linn.). International Journal of Phytopharmacology, 2014; 5(4): 328-334.
8. Eman Mohammed El- Metwally. Evaluation of Antiulcer Activity of Ginger, Clove and Castor oils against Aspirin induced Gastric ulcers in rats. World Applied Sciences Journal, 2014; 29(7): 815- 824.
  9. Rachhadiya Rakesh M, Kabra Mahaveer Prasad, Rajkumar Shete V. Evaluation of Antiulcer activity of Castor oil in Rats. International Journal of Research in Ayurveda and Pharmacy, 2011; 2(4): 1349-1353.
  10. Priyanka Yadav, Aditya Ganeshpurkar, Gopal Rai. *In vitro* H<sup>+</sup> -K<sup>+</sup> ATPase inhibitory potential of methanolic extract of *Cissus quadrangularis* Linn.. Pharmacognosy Research, 2012; 4(2): 123-126.
  11. Samuel Onasanwo A, Neetu Singh, Samuel Olaleye B, Vaibhav Mishra, Gautam Palit. Antiulcer and Antioxidant activities of *Hedranthera barteri* {(Hook F.) Pichon}. International Journal of Medical Research, 2010; 132: 442-449.
  12. Rajesh Kashi Prakash Gupta, Pradeepa, Manunatha Hanumanthappa. In-vitro antioxidant and H<sup>+</sup>, K<sup>+</sup>-ATPase inhibition activities of *Acalypha wilkesiana* foliage extract. Journal of Pharmacy and Bioallied Sciences, 2013; 5(3): 2114-223.
  13. Barisi Inayor N, Omodule Ibraheem. Assessing *Ricinus communis* L.(castor) whole plant part for Phenolics and Saponin constituents for Medicinal and Pharmaceutical Applications. International Journal of Advances in Pharmacy, Biology and Chemistry, 2014; 3(4): 815- 826.
  14. Momoh AO, Oladunmoye MK, Adebolu TT. Evaluation of the Antimicrobial and Phytochemical Properties of Oil from Castor Seeds (*Ricinus communis* Linn). Bulletin of Environment, Pharmacology and Life Sciences, 2012; 1 (10): 21-27.
  15. Raheela Khurshed, Asia Naz, Erum Naz, Huma Sharif, Ghazala Rizwani H. Antibacterial, Antimycelial and Phytochemical Analysis of *Ricinus communis* Linn., *Trigonella foenum graceum* Linn., *Delonix regia*(Bojer ex Hook) Raf. of Pakistan. Romanian Biotechnological Letters, 2012; 17(3): 7237- 7244.
  16. Manpreet Rana, Hitesh Dhamija, Bharat Prashar, Shivani Sharma. *Ricinus communis* L. – A Review. International Journal of Pharm Tech Research, 2012; 4(4): 1706-1711.
  17. Vandita K, Nirali Amin, Khyati P, Monisha K. Effect of Phytochemical constituents of *Ricinus communis*, *Pterocarpus santalinus*, *Terminalia bellerica* on Antibacterial, Antifungal and Cytotoxic Activity. International Journal of Toxicological and Pharmacological Research, 2013; 5(2): 47- 54.
  18. Gamble JS. Flora of the Presidency of Madras. Vol. 2, London :West, Newman and Adlard, 1915; 886-933.
  19. Neelapu Neelima, Naikwadi Gajanam Devidas, Muvvala Sudhakar, Jadhav Kiran V. A Preliminary phytochemical investigation on the leaves of *Solanum xanthocarpum*. International Journal of Research in Ayurveda and Pharmacy, 2011; 2(3): 845- 850.
  20. Pushpalata More, Gulab Rathod, Narayan Pandhure. Phytochemical Analysis and Antibacterial Activity of *Ricinus communis* L.. Research Gate, 2014; 3(1), 49-51.
  21. Harborne JB, Phytochemical Method: A guide to modern techniques of plant analysis, New York: Chapman and Hall, 1983.
  22. Zhang SL, Li H, He X, Zhang RQ, Sun YH, Zhang CF, Wang CZ, Yuan CS. Alkaloids from *Mahonia bealei* possess anti-H<sup>+</sup>/K<sup>+</sup>-ATPase and anti-gastrin effects on pyloric ligation-induced gastric ulcer in rats. Phytomedicine, 2014; 21(11): 1356-1363.
  23. Qiaoyin Zhang, Nianyu Huang, Junzhi Wang, Huajun Luo, Haibo He, Mingruo Ding, Wei-Qiao Deng, Kun Zou. The H<sup>+</sup>/K<sup>+</sup>-ATPase inhibitory activities of Trametenolic acid B from *Trametes lactinea* (Berk.) Pat, and its effects on gastric cancer cells. Fitoterapia, 2013; 89: 210-217.